

# DRUG FACTS

Active ingredient(s)

Purpose

## Use(s)

temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

## Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause stomach bleeding.

The chance is higher if the child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing an NSAID (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

## Do not use

- if the child has ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

## Ask a doctor before use if you have

- problems or serious side effects from taking pain relievers or fever reducers
- stomach problems that last or come back, such as heartburn, upset stomach, or stomach pain
- ulcers
- bleeding problems
- not been drinking fluids
- lost a lot of fluid due to vomiting or diarrhea
- high blood pressure
- heart or kidney disease
- taken a diuretic

## Ask a doctor or pharmacist before use if you are

- taking any other drug containing an NSAID (prescription or nonprescription)
- taking a blood thinning (anticoagulant) or steroid drug
- under a doctor's care for any serious condition
- taking any other drug

## When using this product

- give with food or milk if stomach upset occurs
- long term continuous use may increase the risk of heart attack or stroke

## Stop use and ask a doctor if

- the child feels faint, vomits blood, or has bloody or black stools. These are signs of stomach bleeding.
- stomach pain or upset gets worse or lasts
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

## If pregnant or breast-feeding

## Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center right away.

## Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- do not give longer than 10 days, unless directed by a doctor (see Warnings)
- shake well before using
- find right dose on chart below. If possible, use weight to dose; otherwise use age.
- use only with enclosed measuring cup
- if needed, repeat dose every 6-8 hours
- do not use more than 4 times a day
- replace original bottle cap to maintain child resistance
- wash dosage cup after each use

Dosing Chart		
Weight (lb)	Age(yrs)	Dose (teaspoonful or mL)
under 2 years		ask a doctor
24-35 lbs	2-3 years	1 tsp or 5 mL
36-47 lbs	4-5 years	1 ½ tsp or 7.5 mL
48-59 lbs	6-8 years	2 tsp or 10 mL
60-71 lbs	9-10 years	2 ½ tsp or 12.5 mL
72-95 lbs	11 years	3 tsp or 15 mL

## Other information

- do not use if printed neckband is broken or missing
- store at 20-25°C (68-77°F)

- see bottom panel for lot number and expiration date

***Inactive ingredients***

anhydrous citric acid, butylparaben, corn syrup, D&C red #33, FD&C blue #1, FD&C red #40, flavor, glycerin, hypromellose, polysorbate 80, propylene glycol, purified water, sodium benzoate, sorbitol, xanthan gum